

INTRAVESICAL CHEMOTHERAPY

Why do I need this done ?

If your doctor has arranged for you to have this procedure, this brochure gives you the basic facts on what is going to happen to you and your body, in an easy to understand way. Intravesical chemotherapy using Mitomycin C (MMC) is a procedure that patients have following surgery or biopsy for bladder cancer. It involves exposing the bladder lining to MMC which reduces the chance of your bladder cancer returning following surgical removal.

What to expect before the procedure ?

MMC therapy is typically performed as an outpatient procedure and you will be awake. Prior to the procedure it is important you follow your surgeon/hospital's instructions which may include additional investigations, medication changes and/or a short course of antibiotics. You should contact your treating team if you have symptoms of urine infection prior to the procedure. Upon arrival to the hospital facility, another urine sample will be taken to ensure there is no urinary tract infection. If there is no infection you will be able to proceed with the procedure.

What to expect during the procedure ?

After cleaning the genital area with antiseptic solution, numbing gel is applied to the natural water pipe (the urethra). A catheter is then inserted into the bladder to enable us to deliver the MMC preparation into the bladder. The catheter is then removed and you will be asked to not pass urine for the next hour to allow the medicine the time to work. Your hospital may allow you to leave after this and pass urine at home.

What to expect after surgery

It is normal to experience the following side effects up to 2-3 days after your MMC therapy:

Painful and/or frequent urination

Small amount of blood in urine

Flu-like symptoms such as:

Headache, low-grade fever, poor appetite, low energy, shivers

You will be given specific safety instructions to follow for the next couple of days. It is advisable to avoid sexual intercourse for at least 24 hours after each treatment.

Going home tips

MMC can make you feel tired and under the weather. Take it easy for a couple days


If you are experiencing painful urination, over the counter medication such as Ural can help. Drink plenty of water. It is normal for there to be a small amount of blood in your urine for 2-3 days after the procedure.

Next steps

Typically, you will receive MMC therapy once a week for 6 weeks, which is known as the 'induction' phase. After this time many people move on to a 'maintenance' phase where they receive further doses every 4 weeks for up to a year.

Bladder cancer can recur during or after MMC treatment, so regular follow-up bladder examinations (cystoscopies) are an important part of your follow-up.

COMMON	OCCASIONAL	RARE
Minor side-effects (as mentioned above)	Failure to complete treatment due to bladder discomfort Urethral scarring Urine infection requiring antibiotics	Severe pain on instillation that persists after you have emptied your bladder Allergic reaction to MMC Long-term bladder shrinkage and reduced bladder capacity



Disclaimer: The information in this publication is of a general, non-specific nature. It is not intended, and should not be used as a substitute for the advice of qualified medical professionals.

SEE OUR WEBSITE FOR FREE DOWNLOAD OF OUR BLADDER CANCER CARE PLAN TO FILL OUT WITH YOUR DOCTOR:

WWW.BLADDERCANCER.ORG.AU/BLADDER-CANCER-CARE-PLAN/



BladderCancer.org.au